

AD-A184 985

EVOLUTION OF MEAN SQUARE MAGNETIC VECTOR POTENTIAL IN
COMPRESSIBLE TWO-DI. (U) NAVAL RESEARCH LAB WASHINGTON
DC R B DAHLBURG 08 SEP 87 NRL-MR-6812


1/1

UNCLASSIFIED

F/G 28/9

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963 A

Naval Research Laboratory

Washington, DC 20375-5000

DTIC FILE COPY



NRL Memorandum Report 6012

AD-A184 985

**Evolution of Mean Square Magnetic
Vector Potential in Compressible,
Two-Dimensional, Magnetofluid Turbulence**

R.B. DAHLBURG

*Center for Computational Physics Developments
Laboratory for Computational Physics and Fluid Dynamic*

September 8, 1987

DTIC
ELECTE
SEP 25 1987
S D
E

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Memorandum Report 6012			7a. NAME OF MONITORING ORGANIZATION		
6a. NAME OF PERFORMING ORGANIZATION Naval Research Laboratory		6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code)		
6c. ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	10. SOURCE OF FUNDING NUMBERS		
8c. ADDRESS (City, State, and ZIP Code) Arlington, VA 22217		PROGRAM ELEMENT NO. 61153N	PROJECT NO. RR011-09-43	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Evolution of Mean Square Magnetic Vector Potential in Compressible, Two-Dimensional Magnetofluid Turbulence					
12. PERSONAL AUTHOR(S) Dahlburg, R.B.					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 1987 September 8	15. PAGE COUNT 11
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Compressible Magnetohydrodynamic turbulence		
			Numerical simulation		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) <p>The mean square magnetic vector potential is not invariant in ideal, two dimensional, compressible, magnetohydrodynamic turbulence. A new, related invariant for the compressible case is given. The results are demonstrated by numerical simulation. In the compressible case large amplitude fluctuations in the mean square magnetic vector potential are observed.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL R.B. Dahlburg			22b. TELEPHONE (Include Area Code) 202-767-6326		22c. OFFICE SYMBOL

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

CONTENTS

Introduction	1
Results of numerical simulations	4
Discussion	5
Acknowledgements	6
References	6

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



EVOLUTION OF MEAN SQUARE MAGNETIC VECTOR POTENTIAL IN COMPRESSIBLE, TWO-DIMENSIONAL, MAGNETOFLUID TURBULENCE

Introduction

Incompressible, two-dimensional magnetohydrodynamic (MHD) turbulence has been investigated under a wide range of conditions^{1 2 3 4 5 6 7}. The starting point of these investigations is the identification of the quantities which remain constant in the absence of physical dissipation. A striking difference in compressible, two-dimensional MHD is the absence of the mean square magnetic vector potential as an ideal invariant. In this report we discuss this absence, and suggest a related invariant for the compressible case. The supporting results of numerical simulations are then given.

The nonlinear partial differential equations which govern the behaviour of an ideal, two-dimensional, compressible magnetofluid, written in a dimensionless form, are:

$$\frac{\partial \rho}{\partial t} = -\nabla \cdot (\rho \mathbf{v}), \quad (1a)$$

$$\frac{\partial(\rho \mathbf{v})}{\partial t} = -\nabla \cdot (\rho \mathbf{v} \mathbf{v} - \mathbf{B} \mathbf{B} + \frac{1}{2}(p + B^2)\mathbf{I}), \quad (1b)$$

$$\frac{\partial A}{\partial t} = \mathbf{v} \times \mathbf{B}, \quad (1c)$$

$$\frac{\partial E}{\partial t} = -\nabla \cdot ((E + p)\mathbf{v} + (|\mathbf{B}|^2 \mathbf{I} - 2\mathbf{B}\mathbf{B}) \cdot \mathbf{v}), \quad (1d)$$

where $\rho(x, y, t) \equiv$ mass density, $\mathbf{v}(x, y, t) \equiv$ flow velocity, $p(x, y, t) \equiv$ mechanical pressure, $\mathbf{I} \equiv$ unit dyad, $A(x, y, t) \equiv$ magnetic vector potential, $\mathbf{B}(x, y, t) \equiv$ magnetic induction field, and $E(x, y, t) \equiv$ total energy density. In addition, we utilize an ideal gas equation of state, viz., $p(x, y, t) = (\gamma - 1)U(x, y, t)$, where $U(x, y, t)$ is the internal energy density, and γ is the ratio of specific heats (assumed equal to $\frac{5}{3}$). The normalization⁸ is such that $E_0 = p_0 = B_0^2/8\pi = \rho V_A^2/2$, where V_A^2 is the square of the Alfvén speed. The effects of thermal conduction are not considered.

The ideal conservation properties of the mean square magnetic vector potential are determined by equation 1c¹. Upon multiplying this equation by A , we have:

$$\frac{\partial A^2}{\partial t} = -\nabla \cdot (\mathbf{v}A^2) + A^2 \nabla \cdot \mathbf{v}. \quad (2)$$

Integrating this over a periodic box, or one with perfectly conducting boundary conditions, we have, after some algebra:

$$\frac{\partial}{\partial t} \int A^2 d^2x = \int (A^2 \nabla \cdot \mathbf{v}) d^2x. \quad (3)$$

For an incompressible magnetofluid, $\nabla \cdot \mathbf{v} = 0$, and the right hand side of equation 3 equals zero. In the compressible case, however, the right hand side of this equation can be finite, as can easily be shown by example. Thus, the magnetofluid's compressibility both removes the constraint that the mean square magnetic vector potential be an ideal invariant of the system, and provides a possible source or sink for this quantity.

A new, related, ideal invariant for the compressible, 2-d magnetofluid can be found by utilizing the mass density equation (1a). First, multiply equation 2 by the mass density ρ . After substituting from equation 1a, this gives:

$$\frac{\partial \rho A^2}{\partial t} = -\nabla \cdot (\mathbf{v} \rho A^2). \quad (5)$$

When equation 5 is integrated over a periodic box, the right hand side will equal zero, implying that

$$\int \rho A^2 d^2 x \quad (6)$$

is an ideal constant of the motion for the compressible, two-dimensional magnetofluid. Note that equation 6 reduces to the mean square magnetic vector potential as $\rho \rightarrow 1$. This is the situation in compressible magnetofluids which exhibit negligible variation in mass density.

We note that $\int \rho A^n d^2 x$ is constant for $n = 0, 1, 2, 3, \dots$. Hence the question naturally arises as to why we have singled out the $n = 2$ case for consideration. We believe that this is the significant value of n because of its close analogy with the kinetic energy, $\int \rho |\mathbf{v}|^2 d^2 x$. Both of these quantities exhibit smooth transitions to the incompressible limit, *e.g.*, as $\rho \rightarrow 1$.

Results of numerical simulations

We have written a computer code which solves the physically dissipative version of equations 1a - 1d. For the discretization, a dealiased Fourier pseudospectral method is used which will be described in greater detail elsewhere. For the run described here, 32×32 Fourier modes are used, and $\Delta t = 1/250$. A static, uniform, equilibrium state is considered. To initiate a nontrivial evolution in the magnetofluid, we initialize the magnetic field with random noise. With no physical dissipation, the code should exhibit conservation of the ideal invariants.

Figure 1 shows the evolution of the mean square magnetic vector potential as a function of time. Low-frequency, large amplitude oscillations in this quantity are apparent, suggesting that the compressible term serves alternatively as a source or sink of this quantity. The mean square magnetic vector potential varies by as much as approximately 25 % from its initial value. Figure 2 shows the integral given in equation 6 as a function of time for the same run. Note the difference in scaling between the two figures. The maximum variation of this quantity from its initial value is approximately 0.3 % over the course of the run. The conservation is especially good for the first three Alfvén transit times.

We have repeated these calculations for other randomly initialized cases, with similar results being obtained.

Discussion

Statistical description has proven to be the most useful method for characterizing turbulent magnetofluids. The first step in such descriptions is to identify those quantities which remain invariant for the ideal equations of motion. These invariants serve to define hypersurfaces in the phase space of Fourier coefficients of the independent variables of the system. If such a statistical theory can be formed for compressible, 2-d MHD, then it will differ from the incompressible case from the start because the ideal invariants are not the same.

Absence of conservation of mean square magnetic vector potential in a 2-d compressible, turbulent magnetofluid implies several things. First, an inverse cascade of mean square magnetic vector potential is no longer to be expected, since this is not a conserved quantity. Second, there exists in this case the possibility of a 2-d compressible turbulent dynamo. A related matter is the following: the magnetic energy need no longer decay selectively with respect to the mean square magnetic vector potential⁴, since the decay of this latter quantity can be hastened or retarded by compressibility effects.

The new invariant which we have given reduces to the mean square magnetic vector potential in the limit of weak compressibility, where the variation of the mass density will be insignificant. Perhaps for this reason the mean square magnetic vector potential has proven a useful quantity in compressible magnetofluids like the solar wind, in which there is little variation in the mass density. In other magnetofluids, *e.g.*, the upper solar atmosphere, variation of the mass density cannot be ignored. The new invariant (equation 6) takes the mass density variation of the magnetofluid into account. We conjecture that the new invariant replaces the

mean square magnetic vector potential in compressible, 2-d, magnetofluid turbulence. Further calculations will be required to support this conjecture. The rather difficult question of identifying the regions of Fourier space where the Fourier transformed version of equation 6 remains positive definite must also be addressed (a problem which must also be faced with respect to the kinetic energy). We also note here that the results of this paper can be extended in a straightforward way to the conservation of enstrophy in a compressible, 2-d, neutral fluid which is either isentropic or barotropic.

Acknowledgements

Helpful comments by Dr. J. P. Dahlburg, Dr. J. M. Picone, and Dr. D. C. Montgomery are gratefully acknowledged. This work was sponsored by the National Aeronautics and Space Administration Solar Terrestrial Theory Program and by the Office of Naval Research. The numerical simulations reported here were sponsored by a generous grant of computer time on the NRL CRAY-XMP from the director of research of the Naval Research Laboratory.

References

- ¹D. Fyfe and D. Montgomery, *J. Plasma Phys.* 16, 181 (1976).
- ²S. A. Orszag and C. -M. Tang, *J. Fluid Mech.* 90, 129 (1979).
- ³R. H. Kraichnan and D. Montgomery, *Rep. Prog. Phys.* 43, 547 (1980).
- ⁴W. H. Matthaeus and D. Montgomery, *Ann N.Y. Acad. Sci.* 357, 203 (1980).
- ⁵U. Frisch, A. Pouquet, P. -L. Sulem, and M. Meneguzzi, *J. Mec. Theo. Appl., Numero Special*, 191 (1983).
- ⁶R. B. Dahlburg, T. A. Zang, and D. Montgomery, *J. Fluid Mech.* 169, 71 (1986).
- ⁷W. H. Matthaeus and S. L. Lamkin, *Phys. Fluids* 29, 2513 (1986).
- ⁸D. Schnack and J. Killeen, *J. Comp. Phys.* 35, 110 (1980).




Figure 1. Plot of mean square magnetic vector potential, $\int A^2 d^2x$, vs time.




Figure 2. Plot of $\int \rho A^2 d^2 x$ vs time.

END

11-87

DTIC